





DubMad N	ucleotide Protein Genome Structure PopSet Taxonomy OMIM Book								
PubMed Nu Search PubMed	for Go Clear	J							
Bearen 1 tomos	Limits Preview/Index History Clipboard Details								
Entrez PubMed	Display Abstract  Sort  Save Text  Clip Add  Order								
	1: Curr Opin Immunol 1994 Jun;6(3):407-13 Related Articles, Books, LinkC Full Toodon BloMedNet @ Tumor necrosis factor receptor superfamily members and their	)ut							
5 104 1	ligands.								
PubMed Services	Armitage RJ.								
	Department of Cellular Immunology, Immunex Research and Development Corporation, Seattle, Washington 98101.								
Related Resources	The tumor necrosis factor (TNF) receptor family comprises a number of type I integral membrane glycoproteins which exhibit sequence homology in their cysteine-rich extracellular domains. Recently, ligands for many of these receptors have been identified. These molecules all display sequence identity with TNF and lymphotoxin beta, prototype ligands for this receptor family, and have the conformation of type II transmembrane molecules. While certain biological activities are common to many members of this TNF-like family, other activities appear to be shared only by some ligands, or are unique. The identification of the TNF superfamily of ligands has provided the opportunity to compare and contrast the diverse biological effects mediated through the interaction of these related molecules with their receptors.								
	Publication Types:  Review  Review, Tutorial								
	PMID: 7917108 [PubMed - indexed for MEDLINE]								

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

▼ Save

Text

Clip Add

Order

₹

Sort

Display

Abstract







- NCDI		•	ab y	40,	of Medicine NEW						
PubMed No	ucleotide	Protein	Genom	ie S	Structur	e F	PopSet	Taxonom	y OMIM	Books	
Search PubMed	▼ for _		. /						Go Clear		
	Limits	Prev	iew/Index	His	tory	Clip	board	Details			
	Display	Abstract	₹	Sort	▼	Save	Text	Clip Add	Order	· ·	
Entrez							<u> </u>	<u>.                                    </u>	3		
PubMed		:283-98 RSCIENCE TARTICLE	-				•		ed Articles,	LinkOut	
PubMed Services	prote	ein (MO	-	1CP-2	and	MCF	P-4, mF	onocyte cl RNA expr sts.			
	Benarafa C, Cunningham FM, Hamblin AS, Horohov DW, Collins ME.										
	Department of Pathology and Infectious Diseases, The Royal Veterinary College, Hatfield, Hertfordshire AL9 7TA, UK. benarafa@rvc.ac.uk										
Related Resources	We report the cloning of four equine CC chemokines, eotaxin, monocyte chemoattractant protein (MCP)-1, MCP-2 and MCP-4, which show high levels of identity with their respective homologous sequences in other species. Using a multiplex RT-PCR, we have studied the constitutive mRNA expression of these four CC chemokines in skin, lung, liver, spleen, jejunum, colon and kidney of normal adult horses and compared this data with the eosinophil counts in the same samples. We demonstrate that eotaxin mRNA is only expressed in jejunum and colon, where there are large numbers of eosinophils suggesting that eotaxin might be recruiting eosinophils in the normal digestive tract of the horse. MCP-1 and MCP-4 are expressed in all tissues whereas MCP-2 is only found in some samples of lung, spleen, liver and kidney. We also report the early induction (2h) of equine eotaxin and MCP-4, and the up-regulation of MCP-1 by interleukin-4 in dermal fibroblasts, suggesting these chemokines might be involved in equine skin allergic diseases.  PMID: 11044560 [PubMed - indexed for MEDLINE]										
	Display	Abstract	ন	Sort	[F]	Save	Text	Clip Add	Order		

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer